

APPLICATION

Add 1 lb. of Super K Fulvate into 120 gallons of water
Dissolve rate for foliar spray: 1: 2500-3000 Dissolve rate for root irrigation: 1: 1800-2500

	Foliar Application Times (800-1200gr /acre) Root irrigation (1200-1500gr /acre)			
	First	Second	Third	Optional
Melon, Eggplants, Pepper, Squash, ect	Six leaf stage	Early bloom	First fruit set	4 weeks later
Bean, Peas	Six leaf stage	First bloom	First pods	
Broccoli, Cabbage, Cauliflower	Six true leaf stage	3 weeks later	Head initiation	
Cucumbers, Cucurbits	Six leaf stage	Just prior to bloom	Whilst picking	
Maize, Sweet Corn	Six leaf stage	55-75cm/20-30 inch growth	Just prior to tasseling	
Leek, Onions, Carrot, Turnips	2-3 weeks after emergency	Root enlargement	Every two weeks until harvest	
Potatoes	Six leaf stage	When tuber approx. pea size	Early bloom	Bulking up
Tomato (fresh fruit market)	Six leaf stage	Early bloom	Whilst picking	After 2 weeks

Turf: 800-1200 grams /acre or 150-200 grams/100L of water

Golf Courses
Greens: 800-1200grams /acre

- 1.Recommended to apply "little and often", once or twice a month throughout the year
- 2. Regular applications to greens and tees are especially beneficial
- 3. Applied as bio-stimulant to increase the number and activity of micro-organism. This will improve grass growth and help reduce soil borne disease symptoms in the turf
- 4.Tillering will also be increased but, without excessive top growth



3006 JCS Industrial Drive Brownsville TX 78526 Ph. (956) 574-8280 www.ferti-organic.com